

## Measurements of the CKM angle $\gamma/\phi_3$ at $B$ -factories

YOSHIYUKI ONUKI ON BEHALF OF THE BELLE AND BABAR COLLABORATIONS

*Department of Physics, Tohoku University, 6-3 Aramaki-Aza, Aoba, Sendai, Miyagi,  
980-8578, Japan*

### Abstract

The CKM angle  $\gamma/\phi_3$  have been measured by two B-factories, the PEP-II collider for the BaBar experiment and the KEKB collider for the Belle experiments. The present paper reports recent progress in  $\gamma/\phi_3$ .

PRESENTED AT

The Ninth International Conference on  
Flavor Physics and CP Violation  
(FPCP 2011)  
Maale Hachamisha, Israel, May 23–27, 2011

# 1 Introduction

The  $CP$  violation in the Standard Model is explained by the quark mixing matrix of the charged current weak interaction. The matrix is as known as Cabibbo-Kobayashi-Maskawa(CKM) matrix [1]. The element has the irreducible complex phase which causes  $CP$  violation in  $K$  and  $B$  meson systems [2, 3, 4]. The unitarity of CKM matrix has bring some constrains of the elements  $V_{i,j}$  to draw triangles in the complex plane, where  $i$  and  $j$  stand for the up-type quark ( $u, c, t$ ) and down type quark ( $d, s, b$ ), respectively. One of the triangles which has the sides in same order of magnitude is suitable place to measure the elements precisely. The angles,  $\alpha/\phi_2$ ,  $\beta/\phi_1$  and  $\gamma/\phi_3$  has been measured by two  $B$ -factories, the PEP-II collider for the BaBar experiment [5] and the KEKB collider for the Belle experiments [6], where  $\alpha/\phi_2 = \text{Arg}[-(V_{td}V_{tb}^*)/(V_{ud}V_{ub}^*)]$ ,  $\beta/\phi_1 = \text{Arg}[-(V_{cd}V_{cb}^*)/(V_{td}V_{tb}^*)]$  and  $\gamma/\phi_3 = \text{Arg}[-(V_{ud}V_{ub}^*)/(V_{cd}V_{cb}^*)]$ .

Current most worst precision of the angle is angle  $\gamma/\phi_3$ . Improvement of the precision is eagerly awaited. The  $\gamma/\phi_3$  can be obtained mainly using the process  $B \rightarrow D^{(*)}K^{(*)}$  involved with interference with Cabibbo-suppressed  $b \rightarrow u$  and Cabibbo-favored  $b \rightarrow c$  quark transition [7]. Some methods to extract  $\gamma/\phi_3$  had been suggested so far: GLW [8], ADS [9], Dalitz [10, 11] analyses. The GLW analysis uses  $D$  meson decays to the  $CP$  eigenstates. The ADS analysis uses Cabibbo-favored  $\bar{D}^0$  and doubly Cabibbo-suppressed  $D^0$  meson decays. The Dalitz method uses Cabibbo-allowed three-body  $D$  meson decays.

## 2 GLW analysis and the recent measurements

Gronau, London and Wyler (GLW) [8] proposed analysis method to extract the angle  $\gamma/\phi_3$  with  $D^0$  and  $\bar{D}^0$  decay into  $CP$  eigenstate such as  $K^+K^-$  or  $K_S\pi^0$ , etc. The observables, double ratio and asymmetry, are defined as below.

$$\begin{aligned} R_{CP\pm} &\equiv 2 \frac{\mathcal{B}(B^- \rightarrow D_{CP\pm}K^-) + \mathcal{B}(B^+ \rightarrow D_{CP\pm}K^+)}{\mathcal{B}(B^- \rightarrow D^0K^-) + \mathcal{B}(B^+ \rightarrow \bar{D}^0K^+)} \\ &= 1 + r_B^2 \pm 2r_B \cos \delta_B \cos \phi_3 \end{aligned} \quad (1)$$

$$\begin{aligned} A_{CP\pm} &\equiv \frac{\mathcal{B}(B^- \rightarrow D_{CP\pm}K^-) - \mathcal{B}(B^+ \rightarrow D_{CP\pm}K^+)}{\mathcal{B}(B^- \rightarrow D_{CP\pm}K^-) + \mathcal{B}(B^+ \rightarrow D_{CP\pm}K^+)} \\ &= \pm 2r_B \sin \delta_B \sin \phi_3 / R_{CP\pm} \end{aligned} \quad (2)$$

where the  $D_{CP\pm}$ ,  $r_B$  and  $\delta_B$  represents the  $D$  meson decay into the  $CP$  eigenstate of even(+) and odd(-), ratio of amplitudes between  $B^- \rightarrow \bar{D}^0K^-$  and  $B^- \rightarrow D^0K^-$  defined  $r_B \equiv |A(B^- \rightarrow \bar{D}^0K^-)|/|A(B^- \rightarrow D^0K^-)|$ , the difference of strong phase, respectively.

The  $B^\pm \rightarrow D_{CP\pm}^{(*)}K^{(*)\pm}$  had been measured BaBar [12, 13, 14] and Belle [15] collaboration. The BaBar recently has reported the measurement of  $B^\pm \rightarrow D_{CP\pm}K^\pm$  following  $D_{CP+} \rightarrow K^+K^-$  and  $\pi^+\pi^-$  for  $CP$ -even,  $D_{CP-} \rightarrow K_S\pi^0, K_S\pi^0, K_S\omega$  and  $K_S\phi$  for  $CP$ -odd eigenstate with the full dataset of  $467 \times 10^6 \Upsilon(4S) \rightarrow B\bar{B}$  decays [16]. The results of combined  $D$  sub-decays as for the each  $CP$  eigenstate are as below.

$$\begin{aligned} A_{CP+} &= 0.25 \pm 0.06 \pm 0.02 \\ A_{CP-} &= -0.09 \pm 0.07 \pm 0.02 \\ R_{CP+} &= 1.18 \pm 0.09 \pm 0.05 \\ R_{CP-} &= 1.07 \pm 0.08 \pm 0.04 \end{aligned} \quad (3)$$

where the first error is statistical, the second error is systematic. The parameter  $A_{CP+}$  is different from zero with a significance of  $3.6\sigma$  standard deviations. The result indicates the direct  $CP$  violation followed Dalitz analysis results in the latter Section. Also, they remove the  $D_{CP-} \rightarrow K_S\phi$  subsample with those from the  $B^\pm \rightarrow DK^\pm$ ,  $D \rightarrow K_S h^+ h^-$  to compare the Dalitz analysis [28]. The measured  $A_{CP\pm}$  and  $R_{CP\pm}$  parameters can determine the  $x_\pm$  and  $y_\pm$  defined in Eq. 16 with the relation as below.

$$x_\pm = [R_{CP+}(1 \mp A_{CP+}) - R_{CP-}(1 \mp A_{CP-})]/4 \quad (4)$$

The obtained  $x_\pm$  are consistent with the Dalitz analysis result as below.

$$\begin{aligned} x_+ &= -0.057 \pm 0.039 \pm 0.015, \\ x_- &= 0.132 \pm 0.042 \pm 0.018. \end{aligned} \quad (5)$$

### 3 ADS analysis and the recent measurements

Atwood, Duniety and Soni (ADS) [9] proposed analysis method to extract the angle  $\gamma/\phi_3$  with Cabibbo-favored  $\bar{D}^0$  decay( $CFD$ ) and doubly Cabibbo-suppressed  $D^0$  decay( $DCSD$ ) to adjust the interfering amplitudes have comparable magnitudes through the  $B^- \rightarrow D^{(*)}K^{(*)-}$  decays, where  $D^{(*)}$  means  $D^{(*)0}$  and  $\bar{D}^{(*)0}$ . The final state  $f$  of  $CFD(DCFD)$  from  $D$  meson can be used for ADS such as  $D^0 \rightarrow K^-\pi^+$ ,  $K^-\pi^+\pi^0$  ( $D^0 \rightarrow K^+\pi^-, K^+\pi^-\pi^0$ ), etc. The observables, double ratio and asymmetry, are defined as below.

$$\begin{aligned} R_{ADS} &\equiv \frac{\mathcal{B}(B^- \rightarrow [f]_D K^-) + \mathcal{B}(B^+ \rightarrow [\bar{f}]_D K^+)}{\mathcal{B}(B^- \rightarrow [\bar{f}]_D K^-) + \mathcal{B}(B^+ \rightarrow [f]_D K^+)} \\ &= r_B^2 + r_D^2 + 2r_{BrD} \cos(\delta_B + \delta_D) \cos \phi_3 \end{aligned} \quad (6)$$

$$\begin{aligned} A_{ADS} &\equiv \frac{\mathcal{B}(B^- \rightarrow [f]_D K^-) - \mathcal{B}(B^+ \rightarrow [\bar{f}]_D K^+)}{\mathcal{B}(B^- \rightarrow [\bar{f}]_D K^-) + \mathcal{B}(B^+ \rightarrow [f]_D K^+)} \\ &= 2r_{BrD} \sin(\delta_B + \delta_D) \sin \phi_3 / R_{ADS} \end{aligned} \quad (7)$$

where  $r_D = |A(D^0 \rightarrow f)/A(\overline{D}^0 \rightarrow f)|$  and  $\delta_D$  is strong phase difference between  $\overline{D}^0 \rightarrow f$  and  $D^0 \rightarrow f$ .

The  $B^\pm \rightarrow D^{(*)}K^{(*)\pm}$  had been measured by BaBar [17, 18] and Belle collaboration [19]. The BaBar recently has reported the measurement of  $B^\pm \rightarrow D^{(*)}K^\pm$  following  $D^* \rightarrow D\gamma, D\pi^0$  and  $D \rightarrow K\pi$  with the full dataset of  $467 \times 10^6 \Upsilon(4S) \rightarrow B\overline{B}$  decays [20]. The results are as below.

$$\begin{aligned}
A_{DK} &= -0.86 \pm 0.47 \pm_{-0.16}^{0.12} \\
R_{DK} &= 0.011 \pm 0.006 \pm 0.002 \\
A_{(D\gamma)K}^* &= -0.36 \pm 0.94 \pm_{-0.41}^{0.25} \\
R_{(D\gamma)K}^* &= 0.013 \pm 0.014 \pm 0.008 \\
A_{(D\pi^0)K}^* &= 0.77 \pm 0.35 \pm 0.12 \\
R_{(D\pi^0)K}^* &= 0.018 \pm 0.009 \pm 0.004
\end{aligned} \tag{8}$$

The BaBar also has updated the  $B^\pm \rightarrow DK^\pm$  following  $D \rightarrow K\pi\pi^0$  with dataset of  $474 \times 10^6 \Upsilon(4S) \rightarrow B\overline{B}$  decays [21]. The result is

$$R_{ADS} = (9.1 \pm_{-7.6}^{8.2} \pm_{-3.7}^{1.4}) \times 10^{-3} \tag{9}$$

The Belle also has reported the measurement of  $B^\pm \rightarrow DK^\pm$  following  $D \rightarrow K\pi$  with the full dataset of  $772 \times 10^6 \Upsilon(4S) \rightarrow B\overline{B}$  decays [22]. The results are as below.

$$\begin{aligned}
A_{DK} &= -0.39 \pm_{-0.28}^{0.26} \pm_{-0.03}^{0.04} \\
R_{DK} &= 0.0163 \pm_{-0.0041}^{0.0044} \pm_{-0.0013}^{0.0007}
\end{aligned} \tag{10}$$

The measured  $R_{DK}$  indicates the first evidence of the signal with a significance of  $4.1\sigma$  standard deviations.

## 4 Dalitz analysis and the recent measurements

Dalitz analysis with  $D$  meson decay into  $CP$  eigenstate of three-body decay  $K_S h^+ h^-$  proposed by Giri, Grossman, Soffer, Zupan [10] and Bondar [11] as a effective method to extract the angle  $\gamma/\phi_3$ , where  $h^\pm$  represents charged light hadrons such as pion and kaon. The advantage of the method is only use of Cabibbo-allowed  $D$  decays.

### 4.1 Model-dependent Dalitz analysis and measurements

The BaBar and Belle had reported  $\gamma/\phi_3$  measurement with the model-dependent Dalitz analysis [23, 24, 25]. The model-dependent Dalitz analysis uses the isobar

model [26] which assume three-body decayed  $D$  meson proceed through the intermediate two-body resonances. The total amplitude of Dalitz plane can be represent with two amplitudes of  $D^0$  and  $\overline{D}^0$  decays into same final state of  $K_S h^+ h^-$  as below.

$$f_{B^+} = f_D(m_+^2, m_-^2) + r_B e^{i\phi_3 + i\delta_B} f_D(m_-^2, m_+^2) \quad (11)$$

where  $m_+^2 = m_{K_S h^+}^2$ ,  $m_-^2 = m_{K_S h^-}^2$ . The  $f_D(m_+^2, m_-^2)$  consists of the summed amplitudes of intermediates which comes in the Dalitz plane and single non-resonant amplitude as follows.

$$f_D(m_+^2, m_-^2) = \sum_{j=1}^N a_j e^{i\xi_j} \mathcal{A}_j(m_+^2, m_-^2) + a_{NR} e^{i\xi_{NR}} \quad (12)$$

Where  $a_j$  and  $\xi_j$  are the amplitude and phase of the matrix element,  $\mathcal{A}_j$  is the matrix element of the  $j$ -th resonance, and  $a_{NR}$  and  $\xi_{NR}$  are the amplitude and phase of the non-resonant component.

The Belle had reported the result of  $B^\pm \rightarrow D^{(*)} K^\pm$ ,  $K_S \pi^+ \pi^-$  using 18 two-body amplitudes of isobar model with the dataset of  $657 \times 10^6 \Upsilon(4S) \rightarrow B\overline{B}$  decays [27].

$$\phi_3 = (78.4 \pm_{-11.6}^{10.8} \pm 3.6 \pm 8.9)^\circ \quad (13)$$

where the first error is statistical, the second is systematic, the third error is model uncertainty.

The BaBar has reported the result of  $B^\pm \rightarrow D^{(*)} K^\pm$ ,  $D^* \rightarrow D\pi^0$  and  $D\gamma$ ,  $D \rightarrow K_S \pi^+ \pi^-$  and  $K_S K^+ K^-$  using isobar model with improved 8 two-body amplitudes in each  $D$  meson sub-decays with the dataset of  $468 \times 10^6 \Upsilon(4S) \rightarrow B\overline{B}$  decays [28].

$$\gamma = (68 \pm 14 \pm 4 \pm 3)^\circ \quad (14)$$

where the first error is statistical, the second is systematic, the third error is the model uncertainty.

## 4.2 Model-independent Dalitz analysis and measurements

The model-independent Dalitz analysis had proposed by Giri, Grossman, Soffer and Zupan [10], and further developed by Bonder and Poluektov [29, 30].

Assume that the Dalitz plot is divided into  $2\mathcal{N}$  bins symmetrically to the exchange  $m_-^2 \leftrightarrow m_+^2$ . The bins are denoted by the index  $i$  ranging from  $-\mathcal{N}$  to  $\mathcal{N}$  (excluding 0); the exchange  $m_+^2 \leftrightarrow m_-^2$  corresponds to the exchange  $i \leftrightarrow -i$ . Then the expected number of events in the bins of the Dalitz plot of  $D$  from  $B^+ \rightarrow DK^+$  is

$$N_i^\pm = h_B (K_i + r_B^2 K_{-i} + 2\sqrt{K_i K_{-i}} (x_\pm c_i + y_\pm s_i)) \quad (15)$$

where  $K_i$  is the number of events in the bins in the Dalitz plot of the  $D^0$  in a flavor eigenstate,  $h_B$  is the normalization constant,  $x_{\pm}$  and  $y_{\pm}$  are

$$\begin{aligned} x_{\pm} &= r_B \cos(\delta_B \pm \phi_3), \\ y_{\pm} &= r_B \sin(\delta_B \pm \phi_3). \end{aligned} \quad (16)$$

Coefficients  $c_i$  and  $s_i$ , which include the information about the cosine and sine of the phase difference given by

$$\begin{aligned} c_i &= \frac{\int_{\mathcal{D}_i} \sqrt{p_D \bar{p}_D} \cos(\Delta\delta_D(m_+^2, m_-^2)) d\mathcal{D}}{\sqrt{\int_{\mathcal{D}_i} p_D d\mathcal{D} \int_{\mathcal{D}_i} \bar{p}_D d\mathcal{D}}}, \\ s_i &= \frac{\int_{\mathcal{D}_i} \sqrt{p_D \bar{p}_D} \sin(\Delta\delta_D(m_+^2, m_-^2)) d\mathcal{D}}{\sqrt{\int_{\mathcal{D}_i} p_D d\mathcal{D} \int_{\mathcal{D}_i} \bar{p}_D d\mathcal{D}}}. \end{aligned} \quad (17)$$

These averaged strong phases,  $c_i$  and  $s_i$ , in each bin can be extracted from the quantum-correlated  $D^0$  decays from  $\psi(3770) \rightarrow D\bar{D}$  process. The measurement had been performed by CLEO collaboration [31].

The first model-independent Dalitz analysis of  $B^{\pm} \rightarrow DK^{\pm}$ ,  $D \rightarrow K_S \pi^+ \pi^-$  has been performed by Belle [32]. The results are as below

$$\begin{aligned} x_- &= +0.095 \pm 0.045 \pm 0.014 \pm 0.017, \\ y_- &= +0.137 \pm_{-0.057}^{+0.0053} \pm 0.019 \pm 0.029, \\ x_+ &= -0.110 \pm 0.043 \pm 0.014 \pm 0.016, \\ y_+ &= -0.050 \pm_{-0.055}^{+0.052} \pm 0.011 \pm 0.021, \end{aligned} \quad (18)$$

where the first error is statistical, the second is systematic, the third error is the uncertainty from  $c_i$  and  $s_i$ . The  $(\phi_3, r_B, \delta_B)$  was extracted from  $(x_-, y_-, x_+, y_+)$  using the frequentist treatment with the Feldman-Cousins. The results are

$$\begin{aligned} \phi_3 &= (77.3 \pm_{-14.9}^{+15.1} \pm 4.2 \pm 4.3)^{\circ} \\ r_B &= 0.145 \pm 0.030 \pm 0.011 \pm 0.011 \\ \delta_B &= (129.9 \pm 15.0 \pm 3.9 \pm 4.7)^{\circ}, \end{aligned} \quad (19)$$

where the third error is uncertainty from  $c_i$  and  $s_i$ .

## 5 Conclusion

The measurement precision of  $\gamma/\phi_3$  have been progressed according to the data accumulation at B-factories, and have accelerated by the newly developed efficient physics methods and analysis techniques. Though current statistics of ee colliders is over the 1.2 billion  $B\bar{B}$  pairs, but it is still too small to exploit the  $\gamma/\phi_3$  and the new physics over there. Current most precise determination is brought by the Dalitz analyses.

Both the model-independent and improved model-dependent analysis pushed down the systematic limitation and open up the possibilities of much higher precision determination at super B factories in near future. Furthermore, the recent updated ADS result indicates observation of the Cabibbo-suppressed  $D$  decays which may brought us the competitive determination with the Dalitz analyses. Needless to say, it is important the various approaches including GLW method should be performed since single analysis can't constrain the  $\gamma/\phi_3$  together with the other variables, sufficiently.

## References

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973)
- [2] L. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964)
- [3] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **87**, 091801 (2001); B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **66**, 032003 (2002)
- [4] K. Abe *et al.* (Belle Collaboration), Phys. Rev. Lett. **87**, 091802 (2001); K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66**, 032007 (2002);
- [5] B. Aubert *et al.* (BaBar Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002)
- [6] A. Abashian *et al.* (Belle Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 117 (2002)
- [7] A. B. Carter and A. I. Sanda, Phys. Rev. Lett. **45**, 952 (1980); I. I. Bigi and A. I. Sanda, Phys. Lett. B **211**, 213 (1988).
- [8] M. Gronau, D. London, D. Wyler, Phys. Lett. B **253**, 483 (1991); M. Gronau, D. London, D. Wyler, Phys. Lett. B **265**, 172 (1991);
- [9] D. Atwood, I. Dunietz, A. Soni, Phys. Rev. Lett. **78**, 3257 (1997); Phys. Rev. D **63** 036005 (2001)
- [10] A. Giri, Yu. Grossman, A. Soffer, J. Zupan, Phys. Rev. D **68**, 054018 (2003)
- [11] A. Bondar. Proceedings of BINP Special Analysis Meeting on Dalitz Analysis, 24-26 Sep. 2002, unpublished.
- [12] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **77**, 111102 (2008)

- [13] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **78**, 092002 (2008)
- [14] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **80**, 092001 (2009)
- [15] K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **73**, 051106 (2006)
- [16] P.del Amo Sanchez *et al.* (BaBar Collaboration), Phys. Rev. D **82**, 072004 (2010)
- [17] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **72**, 032004 (2005)
- [18] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **80**, 031102 (2009)
- [19] Y. Horii *et al.* (Belle Collaboration), Phys. Rev. D **78**, 071901(R) (2008)
- [20] P.del Amo Sanchez *et al.* (BaBar Collaboration), Phys. Rev. D **82**, 072006 (2010)
- [21] J. P. Lees *et al.* (BaBar Collaboration), arXiv:1104.4472v1 (2011)
- [22] Y. Horii *et al.* (Belle Collaboration), Phys. Rev. Lett. **106**, 231803 (2011)
- [23] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **95**, 121802 (2005)
- [24] A. Poluektov *et al.* (Belle Collaboration), Phys. Rev. D **73**, 112009 (2006)
- [25] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **78**, 034023 (2008)
- [26] S. Kopp *et al.* (CLEO Collaboration), Phys. Rev. D **63**, 092001 (2001); H. Muramatsu *et al.* (CLEO Collaboration), Phys. Rev. Lett. **89**, 251802 (2002); **90**, 059901 (2003).
- [27] A. Poluektov *et al.* (Belle Collaboration), Phys. Rev. D **81**, 112002 (2010)
- [28] P.del Amo Sanchez *et al.* (BaBar Collaboration), Phys. Rev. Lett. **105**, 121801 (2010)
- [29] A. Bondar and A. Poluektov, Eur. Phys. J. C **47**, 347-353 (2006)
- [30] A. Bondar and A. Poluektov, Eur. Phys. J. C **55**, 51-56 (2008)
- [31] R. A. Briere *et al.* (CLEO Collaboration), Phys. Rev. D **80**, 032002 (2009)
- [32] A. Poluektov *et al.* (Belle Collaboration), Moriond EW 2011 preliminary